

# LONG-TERM WIND SPEED TRENDS IN NORTHWESTERN EUROPE

Neil Atkinson, Keir Harman, Matthew Lynn, Adam Schwarz, Andrew Tindal  
*Garrad Hassan and Partners Ltd.*

*St. Vincent's Works, Silverthorne Lane, Bristol BS2 0QD, UK*

*Tel: +44 (0)117 972 9900 Fax: +44 (0)117 972 9901 email: neil.atkinson@garradhassan.com www.garradhassan.com*

## ABSTRACT

Over recent years the windiness index system has become a widely used tool within the wind energy industry, being used for operational wind farms to understand monthly production, as a reference when estimating future long-term production, and more recently as a tool for assessing portfolio effects [1]. A comparison of several of the main windiness indices for northern Europe has been carried out in order to investigate trends in wind speed over the past 15 years. These windiness indices are only available for the past 15 to 20 years, and due to measurement inconsistencies, reliable wind speed measurements are also typically only available for approximately the past 15 years. Therefore, in the absence of reliable and consistent data for the decades before the start of these indices, several alternative data sources have been investigated, which provide valuable insight into the wind speed trends for this longer term period. Three such data sources, which are available from the beginning of the 20<sup>th</sup> century, are compared to the windiness indices and their use as proxies for wind speed is discussed. From these data sources conclusions are drawn relating to the wind speed trends in northern Europe over the past 40 years and the implications for the use of long-term references, including windiness indices, are discussed.

## 1. WINDINESS INDICES

For the purpose of this investigation, windiness indices from Germany [2], Denmark [3], the Netherlands [4] and the UK [5] have been considered alongside pressure-based data sets derived from measurements and isobar maps. Given the correlation observed in annual windiness between the four countries for which windiness indices are available, it is assumed that the results are broadly representative of much of northwestern Europe. However, it is considered that the results will become less applicable with increasing distance from the countries mentioned above.

Figure 1 shows the main windiness indices from Germany, Denmark, the Netherlands and the UK. The former three are all based on energy production data while that from the UK is based on wind speed data. In order to allow comparison of wind and energy based indices, the indices have been normalised by dividing the annual index anomalies by the standard deviation for the period 1990 to 2005. Also plotted in this figure is the annual average of these indices and the linear regression line fitted to these average values.

It is clear from Figure 1 that there is a reasonable correlation between the windiness indices, with an average coefficient of determination,  $R^2$  value, of 0.75. All the indices support the observation that average wind speeds in northwestern Europe were higher in the early 1990s than they have been for the past 5 to 10 years. There may be concerns about the consistency of the indices over the years. However, as all the indices, derived from somewhat different data sets, show the same broad trend, this increases the confidence in this observation.

From the data plotted in Figure 1, it would be easy to conclude that annual mean wind speeds are dropping due

to some regional effect and to make the further hypothesis that a potential cause may be global warming. A consequence for future wind energy projections would be that we should ramp down future expectations of wind speed, the only question being how rapid the ramp down should be.




Figure 1 – Windiness Indices in Northern Europe

However, an investigation of wind speed trends going back further than 1990 indicates that such an assumption is probably incorrect. In order to investigate the past trend in wind speed we need data going back several decades, covering the period before the start of the windiness indices. An obvious source of data is wind measurements. However, changes to measurement equipment, location or procedure at a measurement mast can result in inconsistencies in the data recorded, and taking these inconsistencies into consideration, it is common to have consistent data extending back no more than 15 years and often less.

## 2. PROXIES FOR WIND SPEED

In the absence of wind speed and wind turbine production data prior to the start of the windiness

indices, several potential sources of data have been considered, which could be used to investigate trends in wind speed in northern Europe going back several decades. It is known that the quality of wind measurements has varied substantially over this period. However, due to the simple, slow changing nature of pressure and the simple instruments used to measure it (the mercury barometer, for example, has been in use since the 17<sup>th</sup> Century, long before the anemometer), pressure measurements are not subject to the same number of variables as wind speed measurements, and as such are likely to be more consistent. These data sets are available from the beginning of the 20<sup>th</sup> century but as it is considered that the certainty of the data increases with time, care should be taken when considering the weight to be put on the earlier years of the data. Although it is not known whether any inconsistencies are present within the data sets, it is considered that as they are based on synoptic charts and sea level pressure measurements, these proxies are likely to be some of the most reliable available. In particular, the Grosswetterlagen and Jenkinson Lamb data sets are based on weather types, and as such are not likely to be sensitive to inconsistencies in recorded pressure data. An investigation of data recorded over the past few decades is deemed sufficient to gain an understanding of possible future wind speed trends and therefore, for the purpose of this investigation, data have been considered from 1965 onwards. The North Atlantic Oscillation (NAO), Grosswetterlagen (GWL) and Jenkinson Lamb indices discussed below are considered to be representative of northwestern Europe, northern Europe and the British Isles respectively.

**2.1. NORTH ATLANTIC OSCILLATION**

A widely used index in meteorology is the North Atlantic Oscillation (NAO) index. This is commonly defined as the difference in normalised pressure between a station in the Azores and one in Iceland. Although the NAO occurs throughout the year, it has a particularly dominant effect on the climate in winter months. In general, a positive NAO index is associated with windier, warmer and wetter winters in northern Europe and a negative NAO index is associated with less windy, colder and drier winters.




Figure 2 – North Atlantic Oscillation Index

A monthly NAO index derived using pressure data from Gibraltar and SW Iceland is available to download from the University of East Anglia Climatic Research Unit [6] for the period from 1821 to the present. To calculate this index, the monthly pressure measurements at each station are normalised by dividing the anomalies by the monthly standard deviation over a defined reference period. The difference between the two normalised pressures is then calculated. Figure 2 shows the NAO index for the period 1965 to 2005, along with the 5 year running average.

**2.2. KATALOG DER GROSSWETTERLAGEN EUROPAS**

Katalog der Grosswetterlagen Europas (GWL) [7], translated as “catalogue of large-scale weather patterns in Europe”, is a taxonomy of large-scale weather patterns over Europe based on a subjective analysis of meteorological charts. For each day from 1881 to the present, the large-scale weather conditions are classified into one of 29 types or as ‘unclassifiable’. The work of Pryor et al [8] has shown that only a few types, which occur frequently, contribute to above average wind speeds over the Baltic region. Namely, westerly anticyclonic, westerly cyclonic and northwesterly cyclonic (Types 1, 2 and 8). According to Pryor et al [8], the GWL classification has been extensively used in previous research and has been demonstrated to be free of artificial bias or trends. A GWL index has been created by taking the total annual occurrence of classification types 1, 2 and 8 and normalising each value to the 1965 to 2004 mean annual occurrence. The GWL index is plotted in Figure 3 for the period 1965 to 2004 along with the 5 year running average.




Figure 3 –Grosswetterlagen Index

**2.3. JENKINSON LAMB WEATHER TYPES**

The Lamb weather classification is a widely used daily large-scale weather classification for the British Isles. The data set was created by subjective analysis of the weather system over the British Isles each day and classification into one of 28 types or as ‘unclassifiable’. An objective scheme to classify the daily weather system according to the Lamb weather types was later developed by Jenkinson and Collison using daily grid point mean sea level pressure data. The data used here are those derived using this objective scheme and are available to

download for the period 1880 to 2004 from the University of East Anglia Climatic Research Unit [9].

In order to ascertain which, if any, of the Lamb weather types are associated with high wind speeds, mean daily wind speed data from four UK Met Office stations for the period 1994 to present were compared with the Jenkinson Lamb weather types data and a mean wind speed associated with each weather type was derived. It was found that Lamb weather types 13 to 17 and 23 to 27 are associated with above average mean daily wind speeds. An index has been created by taking the annual occurrence of these weather types and normalising each value to the 1965 to 2004 mean. The Jenkinson Lamb index is plotted in Figure 4 for the period 1900 to 2004.




Figure 4 –Jenkinson Lamb Index

### 3. DISCUSSION OF PROXIES

In Figure 5, the three proxies for wind speed described above are plotted alongside the average of the four windiness indices (the UK, Denmark, IWET and the Netherlands) for the period 1990 to 2005. To allow comparison, the indices have been normalised by dividing the index anomalies for each annual value by the long-term standard deviation. Correlations of the proxies and the average of the windiness indices indicate an average coefficient of determination,  $R^2$  value, of 0.49. It is clear that, although the correlation quality is not high, on the whole the proxies follow the same trend as the windiness indices and it is considered that the correlations are strong enough to justify the qualitative but not quantitative use of these proxies. It is assumed that the relationship observed between the proxies and the windiness indices over the period 1990 to 2005 would hold were the windiness indices available going back further. It is apparent that the three proxies for wind speed show a similar downward trend to that exhibited by the windiness indices over the period 1990 to 2005 shown in Figure 5.

In Figure 6 the NAO, the GWL, and the Jenkinson Lamb indices are plotted for the period 1965 to 2004 along with the 5 year running average. For comparison purposes, the indices have been normalised by dividing the index anomalies for each value by the standard deviation for the period 1965 to 2004.




Figure 5 –Wind speed proxies and average of windiness indices

The data plotted in Figure 6 provide valuable information on the trend in wind speed prior to the start of the windiness indices discussed in Section 1. It is clear that the recent downward trend does not exist prior to about 1990. Inspection of the NAO, GWL, and Jenkinson Lamb indices for the period since 1900 indicates that there have been several fluctuations in the indices. This suggests that the recent downward trend in mean annual wind speed seen in the indices may represent a return to the longer-term mean, and therefore it would be illogical to conclude that this downward trend in wind speed will continue for the next 15 to 20 years.




Figure 6 – North Atlantic Oscillation, Grosswetterlagen and Indices

It is also implied from the proxies for wind speed discussed here that the period centred on the early 1990s experienced unusually high wind speeds. The data in Figure 6 show that this period, while having higher wind speeds than the period since the mid 1990s also appears to have had higher wind speeds than the previous 20 to 30 years. In fact, inspection of these data sets prior to 1965 indicates that this period was one of the windiest periods, if not the windiest period since the beginning of the century. An investigation carried out by Pryor et al [10] using reanalysis data sets concluded that the 1990s exhibited atypically high wind energy density.

If it were possible to obtain a source of reference wind speed data that were consistent for the past 30 to 40 years, it is considered that the inclusion of this period of

higher windiness would have little effect on the average wind speed for the reference data. However, due to consistency considerations relating to both windiness indices and wind speed measurements, such a long consistent period is not typically available. Long-term references used in energy production assessments typically extend back no more than around 15 years, and as such, are likely to include an unrepresentative proportion of data from this windy period. Where this is the case it is considered that care is needed in the selection of the most appropriate long-term period to use.

#### **4. CONCLUSIONS**

From investigation of several northern European windiness indices and three proxies for wind speed, the following conclusions are drawn regarding long-term trends in wind speed.

- Several northwestern European windiness indices have been obtained and a comparison between them carried out. A similar downward trend in wind speed over the past 15 years can be observed in all of these indices.
- Due to changes in measurement equipment, location and procedure at measurement masts, consistent wind speed data going back more than 15 years are not typically available. The windiness indices compared above are only available since between 1979 and 1989, although the usefulness of the early years is questionable due to the small number of contributing data sources. Therefore, in the absence of reliable and consistent data for the several decades before the start of the wind indices, several alternative data sets have been investigated that cover this period; the North Atlantic Oscillation, Grosswetterlagen and Jenkinson Lamb indices are presented in Sections 2.1 to 2.3.
- It has been demonstrated to an acceptable degree that these three data sets show reasonable agreement with the windiness indices, with an average coefficient of determination,  $R^2$  value, of 0.49. It is therefore considered reasonable to use these data sets as proxies for wind speed prior to the start of the windiness indices. It is clear that, although the correlation quality is not high, on the whole the proxies follow the same trend as the windiness indices and it is considered that the correlations are strong enough to justify the qualitative but not quantitative use of these proxies.
- The recent downward trend observed in the windiness indices is also reflected in the North Atlantic Oscillation, Grosswetterlagen and Jenkinson Lamb data. However, these proxies for wind speed indicate that there was an upward 'blip' in wind speeds centred on the early 1990s, which suggests that this recent downward trend in mean annual wind speed may represent a return to the longer-term mean. It is therefore concluded that a continued ramp down of future wind speeds should not be assumed.
- Given the trends identified within this paper, care needs to be exercised in the selection of the most appropriate long-term reference period to use for a prediction of the long-term mean wind speed at a proposed wind farm site.

#### **REFERENCES**

1. Use of regional wind energy indices to predict long-term wind farm production and to assess portfolio effect, Keir Harmon, Colin Morgan, Garrad Hassan and Partners Ltd., May 2005
2. IWET Windindex, Betreiber-Datenbasis, <http://www.btrdb.de/btrdb/default.htm>
3. Danish Wind Index, Vindstat, [http://www.vindstat.dk/Excel\\_ark/Vindindeks.xls](http://www.vindstat.dk/Excel_ark/Vindindeks.xls)
4. Netherlands Wind Index, Wind Service Holland, <http://home01.wxs.nl/~windsh/windexen.html>
5. UK Wind Index, Garrad Hassan and Partners Ltd.
6. University of East Anglia Climatic Research Unit, North Atlantic Oscillation data, <http://www.cru.uea.ac.uk/cru/data/nao.htm>
7. Katalog der Grosswetterlagen Europas (1881-2004), Potsdam Institute for Climate Impact Research, September 2005
8. Long-term trends in near-surface flow over the Baltic, S C Pryor and R J Barthelmie, January 2003
9. University of East Anglia Climatic Research Unit, Jenkinson Lamb weather types, <http://www.cru.uea.ac.uk/cru/data/lwt.htm>
10. Inter-annual Variability of Wind Indices across Europe, S C Pryor, R J Barthelmie, J T Schoof, December 2005